arXiv:1106.0631v1 [math.NA] 3 Jun 2011

On positivity of principal minors of bivariate Bézier collocation matrix

Gašper Jaklič[a], Tadej Kanduč*[,b]

[a]*FMF and IMFM, University of Ljubljana and PINT, University of Primorska, Jadranska 19, Ljubljana, Slovenia*
[b]*Turboinštitut d.d., Rovšnikova 7, Ljubljana, Slovenia*

Abstract

It is well known that the bivariate polynomial interpolation problem at domain points of a triangle is correct. Thus the corresponding interpolation matrix M is nonsingular. L.L. Schumaker stated the conjecture, that the determinant of M is positive. Furthermore, all its principal minors are conjectured to be positive, too. This result would solve the constrained interpolation problem. In this paper, the basic conjecture for the matrix M, the conjecture on minors of polynomials for degree ≤ 17 and for some particular configurations of domain points are confirmed.

Key words: Collocation matrix, principal minor, bivariate Bernstein polynomial, positivity.
2000 MSC: 65D05, 65F40.

---

1. Introduction

Positivity of determinants (or minors) of collocation matrices is an important property in approximation theory. Nonsingularity of such a matrix implies existence and uniqueness of the solution of the associated interpolation problem. Positivity of principal minors, or even total positivity is used in the proofs of some well-known results, see [1, 2, 3], e.g.

Recently, nonsingularity and principal minors of such matrices for polynomial interpolation at Padua-like points and for interpolation by bivariate

---

*Corresponding author.
Email address: `tadej.kanduc@student.fmf.uni-lj.si` (Tadej Kanduč)

Bézier patches were studied in [4] and [5]. A related problem is a construction or an approximation of Fekete points for a given domain, i.e., the interpolation points, which yield the maximal absolute value of the Vandermonde determinant [6].

It is a well known fact that bivariate Bernstein polynomials $\{B^d_{ijk}\}_{i+j+k=d}$ of degree d form a basis of the space of bivariate polynomials of degree $\leq d$. Let $\mathcal{I}_d = \{(i,j,k) : \ \ i+j+k=d, \ \ i,j,k \in \mathbb{N}\cup\{0\}\}$. For every $\binom{d+2}{2}$ points in a domain, which do not lie on an algebraic hyper-surface of degree $\leq d$, the corresponding interpolation problem is correct ([7]). In particular, the problem is correct for *domain points*

$$\mathcal{D}_{d,T} := \{(i/d, j/d, k/d) : \ \ (i,j,k) \in \mathcal{I}_d\},$$

expressed in barycentric coordinates with respect to a given triangle T. Thus the corresponding interpolation matrix $M := [B^d_\eta(\xi)]_{\xi\in\mathcal{D}_{d,T},\, \eta\in\mathcal{I}_d}$ is nonsingular. In [8], a theorem was stated that by choosing an arbitrary nonempty subset $\mathcal{J} \subset \mathcal{D}_{d,T}$ and the corresponding set of indices Γ, the submatrix $M_\Gamma := [B^d_\eta(\xi)]_{\xi\in\mathcal{J},\, \eta\in\Gamma}$ is nonsingular for all $d \leq 7$, and furthermore, $\det M_\Gamma > 0$. The authors of [8] verified the theorem by computer and provided a proof of nonsingularity of principal matrices for some special configurations of domain points. For more details see [5].

A straightforward way of verifying the conjecture by computing principal minors of M is time consuming due to the exponential growth of the number of subsets that need to be analysed, and cannot be done for $d > 7$ using current computational facilities.

Matrices with positive principal minors are known as P-matrices. Much on them is known, see [9, 10], e.g. Unfortunately, those tools could not be applied for the study of the problem at hand, so a different approach will be used.

In this paper, the conjecture on positivity of determinant of the bivariate Bézier collocation matrix M is confirmed. Furthermore, the result is proven for an arbitrary $\Gamma \subset \mathcal{I}_d$ for $d \leq 17$. Thus the constrained Lagrange interpolation problem has an unique solution. This covers all the cases useful in practice, since it is well known that Lagrange polynomial interpolants of high degrees have undesired properties. Some particular configurations of domain points are analysed. A conjecture for an exact lower bound of $\det M_\Gamma$ is stated.

2. Main results

Let $\boldsymbol{i}$ be a weak 3–composition of an integer d, i.e., $\boldsymbol{i} = (i, j, k)$, such that $|\boldsymbol{i}| := i+j+k = d$ and $i, j, k \in \mathbb{N} \cup \{0\}$. Let $\mathcal{I}_d := \{\boldsymbol{i}\}_{|\boldsymbol{i}|=d}$ be a set of all weak 3–compositions of the integer d. The set $\mathcal{I}_d$ consists of $\binom{d+2}{2}$ compositions.

Let T be a triangle in the plane P. Every point $\boldsymbol{v} \in P$ can be written in barycentric coordinates $\boldsymbol{v} = (u, v, w)$, $u + v + w = 1$, with respect to T. The Bernstein basis polynomials of total degree d in barycentric coordinates are defined as

$$B^d_{\boldsymbol{i}}(\boldsymbol{v}) := B^d_{ijk}(u, v, w) := \binom{d}{\boldsymbol{i}} \boldsymbol{v}^{\boldsymbol{i}} := \frac{d!}{i!j!k!} u^i v^j w^k, \qquad |\boldsymbol{i}| = d.$$

Here the standard multi-index notation and a convention $0^0 = 1$ are used.

Let us denote the subset of all compositions with z zeros by $\mathcal{I}^{(z)}_d \subset \mathcal{I}_d$, $z = 0, 1, 2$. Let $\xi_{\boldsymbol{i}} := \xi_{ijk} := \boldsymbol{i}/d$ be a *domain point* of the triangle T, represented in barycentric coordinates. A domain point $\xi_{\boldsymbol{i}}$ is boundary if at least one of its barycentric coordinates is zero, i.e., $\boldsymbol{i} \in \mathcal{I}^{(1)}_d \cup \mathcal{I}^{(2)}_d$.

We are now ready to present the conjecture, stated in [5], that will be tackled in this paper.

Conjecture 1 ([5]). *For a given triangle T and every nonempty set $\Gamma = \{\boldsymbol{i}_1, \boldsymbol{i}_2, \ldots, \boldsymbol{i}_n\} \subset \mathcal{I}_d$, the matrix*

$$M_\Gamma := [B^d_{\boldsymbol{j}}(\xi_{\boldsymbol{i}})]_{\boldsymbol{i},\boldsymbol{j}\in\Gamma} = \begin{bmatrix} B^d_{\boldsymbol{i}_1}(\xi_{\boldsymbol{i}_1}) & B^d_{\boldsymbol{i}_2}(\xi_{\boldsymbol{i}_1}) & \ldots & B^d_{\boldsymbol{i}_n}(\xi_{\boldsymbol{i}_1}) \\ B^d_{\boldsymbol{i}_1}(\xi_{\boldsymbol{i}_2}) & B^d_{\boldsymbol{i}_2}(\xi_{\boldsymbol{i}_2}) & \ldots & B^d_{\boldsymbol{i}_n}(\xi_{\boldsymbol{i}_2}) \\ \vdots & \vdots & \ddots & \vdots \\ B^d_{\boldsymbol{i}_1}(\xi_{\boldsymbol{i}_n}) & B^d_{\boldsymbol{i}_2}(\xi_{\boldsymbol{i}_n}) & \ldots & B^d_{\boldsymbol{i}_n}(\xi_{\boldsymbol{i}_n}) \end{bmatrix}$$

is nonsingular. Furthermore, $\det M_\Gamma > 0$.

A confirmation of Conjecture 1 would imply the following. Let $\Gamma \subset \mathcal{I}_d$ and let $\mathcal{L}(\{B^d_{\boldsymbol{i}}\}_{\boldsymbol{i}\in\Gamma})$ be the given interpolation space. Then the interpolation problem for the points $\{\xi_{\boldsymbol{i}}\}_{\boldsymbol{i}\in\Gamma}$ in the domain would be correct, i.e., the interpolant would exist and be unique. An example is shown in Fig. 1. The conjecture is important for interpolation with spline functions, since some degrees of freedom are determined through the smoothness conditions and the rest through the interpolation conditions (see [11, 5], e.g.).



Figure 1: Let boundary control points $\{(\xi_{\boldsymbol{i}}, c_{\boldsymbol{i}}) : \boldsymbol{i} \in \mathcal{I}_4^{(1)} \cup \mathcal{I}_4^{(2)},\ c_{\boldsymbol{i}} \in \mathbb{R}\}$ of a Bézier surface p and real values z_{211}, z_{121}, z_{112} be fixed. Conjecture 1 ensures that interior control points $\{(\xi_{\boldsymbol{i}}, c_{\boldsymbol{i}}) : \boldsymbol{i} \in \mathcal{I}_4^{(0)},\ c_{\boldsymbol{i}} \in \mathbb{R}\}$ can be set in such a way that $p(\xi_{\boldsymbol{i}}) = z_{\boldsymbol{i}}$, $\boldsymbol{i} \in \mathcal{I}_4^{(0)}$.

Note that the matrix M_Γ is not symmetric. The determinant of M_Γ is independent of the ordering of elements of Γ as long as the same ordering for rows and columns is used. It is common to use the counter-lexicographical ordering $\succ_{\text{c-lex}}$,

$$(d, 0, 0),\ (d-1, 1, 0),\ (d-1, 0, 1),\ (d-2, 2, 0), \ldots,\ (0, 0, d),$$

but a particular ordering of elements in $\mathcal{I}_d$, which yields a block lower triangular matrix $M_{\mathcal{I}_d}$ (see [12]), will be more convenient. A linear ordering $\succ_b$ is defined as: $\boldsymbol{i} \succ_b \boldsymbol{j}$ if one of the following holds true:

1. $\boldsymbol{i} \in \mathcal{I}_d^{(z_1)}$ and $\boldsymbol{j} \in \mathcal{I}_d^{(z_2)}$ for $z_1 > z_2,\ z_i \in \{0, 1, 2\}$,
2. $\boldsymbol{i}, \boldsymbol{j} \in \mathcal{I}_d^{(z)}$ for $z \in \{0, 1, 2\}$ and $\operatorname{sgn}(\boldsymbol{i}) \succ_{\text{c-lex}} \operatorname{sgn}(\boldsymbol{j})$,
3. $\boldsymbol{i}, \boldsymbol{j} \in \mathcal{I}_d^{(z)}$ for $z \in \{0, 1, 2\}$ and $\operatorname{sgn}(\boldsymbol{i}) = \operatorname{sgn}(\boldsymbol{j})$ and $\boldsymbol{i} \succ_{\text{c-lex}} \boldsymbol{j}$.

Here $\operatorname{sgn}(\boldsymbol{i}) = \operatorname{sgn}(i, j, k) := (\operatorname{sgn}(i), \operatorname{sgn}(j), \operatorname{sgn}(k))$. The ordering $\succ_b$ implies that the matrix $M_{\mathcal{I}_d}$ has a structure

$$M_{\mathcal{I}_d} = \begin{bmatrix} M_{\mathcal{I}_d^{(2)}} & & \\ * & M_{\mathcal{I}_d^{(1)}} & \\ * & * & M_{\mathcal{I}_d^{(0)}} \end{bmatrix},$$

where

$$M_{\mathcal{I}_d^{(2)}} = \begin{bmatrix} M_{\{(d,0,0)\}} & & \\ & M_{\{(d,0,0)\}} & \\ & & M_{\{(d,0,0)\}} \end{bmatrix},$$

$$M_{\mathcal{I}_d^{(1)}} = \begin{bmatrix} M_\epsilon & & \\ & M_\epsilon & \\ & & M_\epsilon \end{bmatrix},$$

and $\epsilon := \{\xi_{ij0} \in \mathcal{I}_d : \ i, j \geq 1\}$.

Thus the problem of verifying the positivity of principal minors of the matrix $M_{\mathcal{I}_d}$ is reduced to each diagonal block matrix separately. The matrix M_ϵ is an univariate Bézier collocation matrix and by [13] it is totally nonnegative with positive principal minors. Therefore, the problem reduces to the study of a $\binom{d-1}{2} \times \binom{d-1}{2}$ matrix $M_{\mathcal{I}_d^{(0)}}$, which, unfortunately, represents a very large part of the matrix $M_{\mathcal{I}_d}$ for a large d.

Let us simplify the considered matrix. Let us construct a matrix N_Γ from M_Γ in the following way:

- for every column, divide each element, that corresponds to a polynomial $B^d_{\boldsymbol{i}}$, by $\binom{d}{\boldsymbol{i}}$,
- multiply the obtained matrix by d^d.

Thus an element $B^d_{\boldsymbol{j}}(\xi_{\boldsymbol{i}})$ is transformed into $\boldsymbol{i}^{\boldsymbol{j}}$. Therefore

$$\det N_\Gamma = \frac{d^{d\cdot|\Gamma|}}{\prod\limits_{\boldsymbol{i}\in\Gamma} \binom{d}{\boldsymbol{i}}} \det M_\Gamma. \tag{1}$$

Clearly, the matrix N_Γ is a principal submatrix of $N_{\mathcal{I}_d}$. Since $\operatorname{sgn}\det M_\Gamma = \operatorname{sgn}\det N_\Gamma$, Conjecture 1 holds for M_Γ iff it holds for N_Γ.

The matrix N_Γ has some nice properties. It consists only of non-negative integers, thus determinant computations are exact. The matrix $N_{\mathcal{I}_d}$ has a simpler structure than $M_{\mathcal{I}_d}$ and is closely related to combinatorial objects. Therefore some properties of the matrix $M_{\mathcal{I}_d}$ will be proven via $N_{\mathcal{I}_d}$. Note that some of the properties are not preserved by the transformation $M_\Gamma \to N_\Gamma$.

As an example, diagonal blocks of the matrix $M_{\mathcal{I}_4}$ and the corresponding matrix $N_{\mathcal{I}_4}$ are shown in Table 1.

$\Gamma \subset I_4$	M_Γ	N_Γ
$\{(4,0,0)\}$	1	256
$\{(0,4,0)\}$		
$\{(0,0,4)\}$		
$\{(3,1,0),(2,2,0),(1,3,0)\}$	$\frac{1}{128}\begin{bmatrix} 54 & 27 & 6 \\ 32 & 48 & 32 \\ 6 & 27 & 54 \end{bmatrix}$	$\begin{bmatrix} 27 & 9 & 3 \\ 16 & 16 & 16 \\ 3 & 9 & 27 \end{bmatrix}$
$\{(3,0,1),(2,0,2),(1,0,3)\}$		
$\{(0,3,1),(0,2,2),(0,1,3)\}$		
$\{(2,1,1)\}$	$\frac{1}{32}\begin{bmatrix} 6 & 3 & 3 \\ 3 & 6 & 3 \\ 3 & 3 & 6 \end{bmatrix}$	$\begin{bmatrix} 4 & 2 & 2 \\ 2 & 4 & 2 \\ 2 & 2 & 4 \end{bmatrix}$
$\{(1,2,1)\}$		
$\{(1,1,2)\}$		

Table 1: Diagonal blocks of the matrices $M_{\mathcal{I}_4}$ and $N_{\mathcal{I}_4}$ by using the ordering $\succ_b$.

Now we are ready to present one of the main results of the paper.

Theorem 2. *Let $d \le 17$. Then Conjecture 1 holds true, i.e.,* $\det M_\Gamma > 0$ *for every nonempty subset $\Gamma \subset \mathcal{I}_d$.*

Proof. Fix d, $1 \le d \le 16$. The matrix $M_{\mathcal{I}_d} + M_{\mathcal{I}_d}^T$ is symmetric and positive definite since its Cholesky decomposition exists. Therefore the matrix $M_{\mathcal{I}_d}$ is positive definite, i.e., $\boldsymbol{x}^T M_{\mathcal{I}_d} \boldsymbol{x} > 0$ for all $\boldsymbol{x} \ne \mathbf{0}$. Thus all principal submatrices of $M_{\mathcal{I}_d}$ are positive definite too and all principal minors are positive.

For $d = 17$, the matrix $M_{\mathcal{I}_d} + M_{\mathcal{I}_d}^T$ has three negative eigenvalues. The matrices $M_{\mathcal{I}_d^{(2)}}$ and $M_{\mathcal{I}_d^{(1)}}$ are P-matrices, thus the problem reduces to the study of the matrix $M_{\mathcal{I}_d^{(0)}}$. Since the Cholesky decomposition of the matrix $M_{\mathcal{I}_d^{(0)}} + M_{\mathcal{I}_d^{(0)}}^T$ exists, the rest of the proof is similar to the first part. $\square$

For $d = 18$, the smallest eigenvalue of $M_{\mathcal{I}_d} + M_{\mathcal{I}_d}^T$ is approximately -1.1×10^{-7}. For $d \geq 18$, the number of negative eigenvalues of the matrix $M_{\mathcal{I}_d} + M_{\mathcal{I}_d}^T$ increases with d. Therefore, this approach can not be used to prove the conjecture in general. However, our result covers all the cases, important in practice, since only Lagrange polynomial interpolants of low degrees are useful.

Theorem 2 implies the following result.

Theorem 3. *Let $\Gamma \subset \mathcal{I}_d$ and let $d \leq 17$. Then for any $\{z_{\boldsymbol{i}}\}_{\boldsymbol{i}\in\Gamma}$, there is an unique polynomial of the form*

$$p := \sum_{\boldsymbol{i}\in\Gamma} c_{\boldsymbol{i}} B_{\boldsymbol{i}}^d$$

such that

$$p(\xi_{\boldsymbol{i}}) = z_{\boldsymbol{i}}, \qquad \boldsymbol{i} \in \Gamma.$$

Remark 1. Theorem 3 generalizes [8, Thm. 3]. Its proof avoids computation of all subdeterminants, as was the case in [8].

Remark 2. Note that the matrix $N_{\mathcal{I}_d}$ is positive definite only for $d \leq 4$.

Remark 3. The approach for proving Theorem 2 can also be used to prove a similar result for the trivariate case: the trivariate Bézier collocation matrix for $d \leq 16$ is positive definite, and thus the constrained polynomial Lagrange interpolation problem is unisolvent.

Now let us consider some particular configurations of domain points (and the corresponding choices of Γ) for d arbitrary.

Theorem 4. *Let d be arbitrary and let Γ satisfy one of the following assumptions:*

1. $|\Gamma| \leq 2$,
2. *let one of the components of (i, j, k) be fixed for all elements in Γ,*
3. $\Gamma = \mathcal{I}_d$,
4. $\Gamma = \mathcal{I}_d^{(0)}$,
5. $\Gamma \subset \mathcal{I}_d^{(2)} \cup \mathcal{I}_d^{(1)}$,
6. $\Gamma = \Gamma_1 \cup \Gamma_2$, *where Γ_1 is one of the sets, defined in 1., 2. or 4., and Γ_2 is a set in 5.*

Then $\det M_\Gamma > 0$.

Proof. Let $|\Gamma| \leq 2$. For $|\Gamma| = 1$, the matrix M_Γ is a positive number.

Now let $\Gamma = \{\boldsymbol{i}_1, \boldsymbol{i}_2\}$. Since the largest element of every column in M_Γ is on the diagonal of M_Γ,

$$\det M_\Gamma = \begin{vmatrix} B^d_{\boldsymbol{i}_1}(\xi_{\boldsymbol{i}_1}) & B^d_{\boldsymbol{i}_2}(\xi_{\boldsymbol{i}_1}) \\ B^d_{\boldsymbol{i}_1}(\xi_{\boldsymbol{i}_2}) & B^d_{\boldsymbol{i}_2}(\xi_{\boldsymbol{i}_2}) \end{vmatrix} > 0.$$

Let one of the components of (i, j, k) be fixed. Without loss of generality we may assume that $\boldsymbol{i}_\ell = (i_\ell, j_\ell, k)$, $i_\ell + j_\ell + k = d$, $\ell \in \{1, 2, \ldots, |\Gamma|\}$. By dividing each element of N_Γ by k^k and multiplying each column by a proper constant, the matrix N_Γ transforms to an univariate Bézier collocation matrix, which is a P-matrix by [13].

Now let us consider the case $\Gamma = \mathcal{I}_d$. It follows from [12] and (1) that

$$\det M_{\mathcal{I}_d} = d^{-d\binom{d+2}{2}} \prod_{\boldsymbol{i}\in\mathcal{I}_d} \binom{d}{\boldsymbol{i}} \prod_{k=1}^{\min\{d,3\}} \left(d^{\binom{d-1}{k}} \prod_{i=1}^{d-k+1} i^{(n-i+1)\binom{d-i-1}{k-2}} \right)^{\binom{3}{k}} > 0. \tag{2}$$

Positivity of $\det M_{\mathcal{I}_d^{(0)}}$ follows from

$$\det\, M_{\mathcal{I}_d} = \det M_{\mathcal{I}_d^{(2)}}\, \det M_{\mathcal{I}_d^{(1)}}\, \det M_{\mathcal{I}_d^{(0)}} > 0.$$

For $\Gamma \subset \mathcal{I}_d^{(2)} \cup \mathcal{I}_d^{(1)}$ and $\Gamma = \Gamma_1 \cup \Gamma_2$, the result follows straightforwardly from the structure of the matrix M_Γ. $\square$

Remark 4. The set Γ in Theorem 4(2) corresponds to domain points in the triangle T, lying on a line parallel to some edge of T.

The nonsingularity of the matrix $M_{\mathcal{I}_d}$ follows from the fact that Bernstein polynomials form a basis of the space of bivariate polynomials of degree $\leq d$. Although the proof that $\det M_{\mathcal{I}_d} > 0$ may seem easy, most of the paper [12] is dedicated to the derivation of determinant formula in a closed form (2).

The subset of decompositions Γ in Theorem 4(5) corresponds to the interpolation problem at boundary domain points of the triangle T.

3. Concluding remarks

In the rest of the paper we will state some conjectures which expand the Conjecture 1, and examine some interesting properties of considered matrices.

Conjecture 5. *For d fixed,*

$$\min_{\substack{\Gamma\subset\mathcal{I}_d\\ \Gamma\neq\emptyset}} \det M_\Gamma = \det M_{\mathcal{I}_d},$$

where $\det M_{\mathcal{I}_d}$ *is given in* (2).

Conjecture 6. *For $\ell\in\mathbb{N}$ let*

$$n_d := \begin{cases} \ell^{3\ell}, & d = 3\ell \\ (\ell+1)^{\ell+1}\ell^{2\ell}, & d = 3\ell+1 \\ (\ell+1)^{2\ell+2}\ell^{\ell}, & d = 3\ell+2 \end{cases}.$$

Then

$$\min_{\substack{\Gamma\subset\mathcal{I}_d\\ \Gamma\neq\emptyset}} \det N_\Gamma = n_d.$$

Conjectures 5 and 6 were verified by a computer for $d\le 7$. Let us prove the latter conjecture for $|\Gamma|\le 2$ and arbitrary d. We will need the following lemma.

Lemma 7. *Let $\boldsymbol{x}=(x,y,z)\in\mathbb{R}^3$ and fix $\boldsymbol{i}=(i,j,k)\in\mathcal{I}_d$. Let the function $f(\boldsymbol{x}) := \boldsymbol{x}^{\boldsymbol{i}}$ be defined on*

$$\Omega = \{(x,y,z):\ \ x+y+z=d,\, 0\le x\le d,\, 0\le y\le d-x\}. \qquad (3)$$

Then f has an unique maximum at $\boldsymbol{i}$ and

$$\max_{\boldsymbol{x}\in\Omega} f(\boldsymbol{x}) = \boldsymbol{i}^{\boldsymbol{i}}.$$

Proof. Let $T=\langle(0,0),(d,0),(0,d)\rangle$ be a triangle in the domain and let us define Bernstein polynomial $B_{\boldsymbol{i}}^d$ on T. Since $B_{\boldsymbol{i}}^d = \binom{d}{\boldsymbol{i}}/d^d f$ and $B_{\boldsymbol{i}}^d$ has an unique maximum in the barycentric point $\xi_{\boldsymbol{i}}$, the proof is complete. $\square$

Proposition 8. *Conjecture 6 holds true for $|\Gamma|\le 2$.*

Proof. Let $|\Gamma| = 1$ and let $g(\boldsymbol{x}) := g(x, y, z) := \boldsymbol{x}^{\boldsymbol{x}}$ be a function, defined on the domain Ω as in (3). We are looking for

$$\alpha := \min_{\boldsymbol{x} \in \Omega \cap \mathbb{Z}^3} g(\boldsymbol{x}).$$

An unique local minimum of g in the interior of Ω is obtained as a solution of the normal system $\partial g/\partial x = 0$, $\partial g/\partial y = 0$, and it is reached at $(d/3, d/3, d/3)$. This is a global minimum since $g(d/3, d/3, d/3) < g(\boldsymbol{x})$ for all $\boldsymbol{x}$ at the boundary of Ω.

If $d \equiv 0 \pmod 3$, then $\alpha = (d/3)^d$.

Let us examine the case $d \equiv 1 (\mathrm{mod}\ 3)$. Then $d = (\ell + 1) + \ell + \ell$ for $\ell \in \mathbb{N} \cup \{0\}$. By the symmetry of the function g and since at least one component of $\boldsymbol{x} \in \Omega \cap \mathbb{Z}^3$ is greater or equal to $\ell + 1$, it is enough to consider the case $\ell + 1 \leq x$ only.

Let us define

$$\Omega_x := \{\boldsymbol{x} = (x, y, z) \in \Omega : \ \ell + 1 \leq x\}.$$

Since g has no extreme point in Ω_x, the minimum value is reached at the boundary of Ω_x. Then the minimum is $\alpha = (\ell + 1)^{\ell+1} \ell^{2\ell}$ and it is achieved at $(\ell + 1, \ell, \ell)$. For $\ell + 1 \leq y$ and $\ell + 1 \leq z$, the derivation is analogous.

The case $d \equiv 2 \pmod 3$ is similar to the previous one. Since $N_{\{\boldsymbol{i}\}} = g(\boldsymbol{i})$, $\boldsymbol{i} \in \mathcal{I}_d$, and $n_d = \alpha$, the conjecture for $|\Gamma| = 1$ is proven.

Now let us consider the case $|\Gamma| = 2$. Let us show that

$$\boldsymbol{i}_1^{\boldsymbol{i}_1} = \det N_{\{\boldsymbol{i}_1\}} \leq \det N_{\{\boldsymbol{i}_1, \boldsymbol{i}_2\}} = \boldsymbol{i}_1^{\boldsymbol{i}_1} \boldsymbol{i}_2^{\boldsymbol{i}_2} - \boldsymbol{i}_2^{\boldsymbol{i}_1} \boldsymbol{i}_1^{\boldsymbol{i}_2}$$

for every $\boldsymbol{i}_1$, $\boldsymbol{i}_2 \in \mathcal{I}_d$, $\boldsymbol{i}_1 \neq \boldsymbol{i}_2$.

By Lemma 7 it follows that $\boldsymbol{i}_2^{\boldsymbol{i}_1} < \boldsymbol{i}_1^{\boldsymbol{i}_1}$ and $\boldsymbol{i}_1^{\boldsymbol{i}_2} \leq \boldsymbol{i}_2^{\boldsymbol{i}_2} - 1$, thus

$$0 \leq \boldsymbol{i}_1^{\boldsymbol{i}_1} (\boldsymbol{i}_2^{\boldsymbol{i}_2} - 1) - \boldsymbol{i}_2^{\boldsymbol{i}_1} \boldsymbol{i}_1^{\boldsymbol{i}_2} = \det N_{\{\boldsymbol{i}_1, \boldsymbol{i}_2\}} - \det N_{\{\boldsymbol{i}_1\}}.$$

□

Let us denote an eigenvalue of the matrix $M_{\mathcal{I}_d}$ with index i and multiplicity n by $\lambda_{i,n}^{<d>}$. An interesting interlacing of eigenvalues can be observed.

Conjecture 9. *The spectrum of the matrix $M_{\mathcal{I}_d}$ consists of d different real eigenvalues*

$$1 = \lambda_{1,3}^{<d>} > \lambda_{2,3}^{<d>} > \lambda_{3,4}^{<d>} > \lambda_{4,5}^{<d>} > \cdots > \lambda_{d,d+1}^{<d>} > 0.$$



Figure 2: Eigenvalue interlacing for $M_{\mathcal{I}_d}$, $d \leq 20$.

The eigenvalues satisfy a criterion, similar as in Cauchy interlacing theorem, i.e.,

$$\lambda_{i,*}^{<d+1>} \geq \lambda_{i,*}^{<d>} \geq \lambda_{i+1,*}^{<d+1>}, \qquad i = 1, 2, \ldots, d.$$

The conjecture was verified by a computer for $d \leq 20$. Note that positivity of eigenvalues would follow, if one would prove that $M_{\mathcal{I}_d}$ is a P-matrix.

As an example, the spectra of $M_{\mathcal{I}_{10}}$ and $M_{\mathcal{I}_{10}} + M_{\mathcal{I}_{10}}^T$ are shown in Table 2 and 3, respectively. The interlacing property for $d \leq 20$ can be observed in Fig. 2.

4. Acknowledgements

The first author thanks ARRS for a grant P1-294. The second author thanks TIA for the financial support.

Eigenvalue	Multiplicity
1	3
0.9000	3
0.7200	4
0.5040	5
0.3024	6
0.1512	7
0.06048	8
0.01814	9
3.629×10^{-3}	10
3.629×10^{-4}	11

Table 2: Spectrum of the matrix $M_{\mathcal{I}_{10}}$.

Eigenvalue	Mult.	Eigenvalue	Mult.	Eigenvalue	Mult.
2.341	1	0.2854	1	7.224×10^{-3}	1
2.308	2	0.2675	1	7.038×10^{-3}	2
1.856	2	0.2585	1	6.565×10^{-3}	1
1.841	1	0.2568	2	6.560×10^{-3}	1
1.475	1	0.1225	1	6.399×10^{-3}	2
1.412	2	0.1179	2	6.220×10^{-3}	2
1.303	1	0.1076	2	6.209×10^{-3}	1
1.001	2	0.1033	2	6.752×10^{-4}	2
0.9377	1	0.1028	1	6.600×10^{-4}	2
0.8855	2	0.03647	2	6.574×10^{-4}	1
0.6082	1	0.03531	1	6.160×10^{-4}	2
0.5742	2	0.03253	2	6.079×10^{-4}	1
0.5221	2	0.03241	1	5.951×10^{-4}	1
0.5180	1	0.03126	1	5.923×10^{-4}	1
0.2995	2	0.03115	2		

Table 3: Spectrum of the matrix $M_{\mathcal{I}_{10}} + M_{\mathcal{I}_{10}}^T$ together with eigenvalue multiplicities.

References

[1] C. de Boor, A practical guide to splines, Springer, 1994.

[2] M. Gasca, Spline functions and total positivity, Revista Mathematica 9, 1996.

[3] M. Gasca, C.A. Micchelli, Total positivity and its applications, Springer, 1996.

[4] L. Bos, S. De Marchi, S. Waldron, On the Vandermonde determinant of Padua-like points, Dolomites Research Notes on Approximation, vol. 2 (2009), 1–15.

[5] M.-J. Lai, L.L. Schumaker, Spline functions on triangulations, Cambridge University Press, 2007.

[6] L. Bos, S. De Marchi, A. Sommariva, M. Vianello, Computing multivariate Fekete and Leja points by numerical linear algebra, SIAM J. Numer. Anal. 48 (2010), 1984–1999.

[7] G. Jaklič, J. Kozak, M. Krajnc, V. Vitrih, E. Žagar. Three-pencil lattices on triangulations, Numer. Algorithms 45 (2007), 49–60.

[8] G. Nürnberger, V. Rayevskaya, L.L. Schumaker, F. Zeilfelder, Local Lagrange interpolation with C^2 splines of degree seven on triangulations, in Advances in Constructive Approximation: Vanderbilt 2003, M. Neamtu and E. Saff (eds.), Nashboro Press, Brentwood, (2004) 345–370.

[9] K.G. Murty, On a characterization of P-matrices, SIAM J. Applied Mathematics, 20 (3), (1971), 378–384.

[10] G.E. Coxson, The P-matrix problem is co-NP-complete, Math. Programming, 64(2) (1994), 173–178.

[11] G. Jaklič, J. Modic, On properties of cell matrices, Appl. Math. Comput., 216 (2010), 2016–2023.

[12] J.M. Brunat, A. Montes, The power-compositions determinant and its application to global optimization, SIAM J. Matrix Anal. Appl., 23(2) (2001), 459–471.

[13] C. de Boor, R. deVore, A geometric proof of total positivity for spline interpolation, Math. Comp. 45(172) (1985), 497–504.